

HYPOTHESIS

Integrated saliva-mediated self-cleaning and wound-healing mechanisms in animals: a hypothesis

Tauqeer Ahmad¹ and Murad Ali²

¹Nuclear Institute for Food and Agriculture (NIFA), Peshawar, Pakistan.

²Department of Food Science and Technology, University of Agriculture, Peshawar, Pakistan.

Abstract

Background: Animals are continuously exposed to environmental contaminants, microbial agents, ectoparasites, and minor injuries, yet many species maintain hygiene and recover from superficial wounds without external intervention.

Objective: This hypothesis manuscript proposes an integrated innate defense model in which grooming behavior, saliva-derived antimicrobial and wound-repair factors, tear antimicrobial activity, and sebaceous secretions act together to support self-cleaning and tissue repair.

Proposed mechanism: Animal saliva contains bioactive components, including lysozyme, lactoferrin, histatins, immunoglobulins, and growth factors, that may contribute to microbial control and wound healing when animals lick their skin, fur, or wounds. Tear secretions similarly contain antimicrobial proteins that protect the ocular surface and assist in debris clearance, whereas grooming behavior distributes saliva and skin lipids while mechanically removing dirt and parasites.

Hypothesis: These interacting behavioral, biochemical, and physiological pathways may be especially active during rest periods, when animals are undisturbed and able to groom. The proposed framework suggests that saliva-mediated grooming, tear antimicrobial activity, and skin secretions constitute a coordinated, evolutionarily conserved innate defense system that contributes to hygiene maintenance, infection prevention, and wound repair.

Research needs: Further comparative and mechanistic studies are needed to test this model and to explore whether saliva-derived antimicrobial and healing compounds can inform bio-inspired veterinary and biomedical therapeutic strategies.

Keywords: Animal grooming; Saliva; Wound healing; Innate immunity; Antimicrobial peptides; Tear secretions; Self-cleaning behavior

*Correspondence: toqeersaqi2853@gmail.com

All rights reserved 501/2026

Copyright © 2026 Bangladesh Society for Veterinary Medicine. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Introduction

Animals in natural environments are continuously exposed to dust, pathogens, parasites, and minor injuries. Despite these challenges, many animals maintain hygiene and recover from superficial wounds without external intervention (Scavizzi et al., 2022). Field and clinical observations often note that animals exposed to contaminants may appear cleaner after periods of rest, consistent with the possibility that behavioral and biochemical self-cleaning mechanisms act together over time. This observation provides the conceptual basis for evaluating natural self-cleaning and wound-healing pathways as an integrated innate defense system (Haick & Zheng, 2025).

Self-grooming is widely documented across animal species, including mammals, birds, and rodents. Cats, dogs, rodents, and livestock frequently lick or groom their fur, skin, and wounds (Tynes, 2025). Grooming can support hygiene maintenance, parasite removal, fur and skin conditioning, thermoregulation, and wound care (Hart & Hart, 2018). Biological fluids such as saliva and tear secretions also contain antimicrobial compounds and healing factors that may contribute to these processes (Mudgil, 2022; Vila et al., 2019).

Saliva is a biologically active fluid containing antimicrobial enzymes, immunological components, and growth-promoting factors (Tamai & Kiyoura, 2025). Tear fluid contains antimicrobial proteins that protect ocular surfaces from infection and help remove debris (Eshac et al., 2021). Together with grooming behavior and skin secretions, these pathways may form an integrated natural hygiene system in animals. This article therefore proposes that animals use saliva-mediated grooming, tear antimicrobial activity, and skin protective secretions to maintain hygiene and promote wound healing, particularly during rest or overnight periods.

Saliva Composition and Biological Functions in Animals

Antimicrobial Components of Animal Saliva

Animal saliva is a complex biological fluid composed of enzymes, peptides, proteins, and immune molecules. Several antimicrobial components have been identified in saliva, including lysozyme, lactoferrin, peroxidases, defensins, and immunoglobulins (Maddu, 2019). Lysozyme can

damage bacterial cell walls and thereby reduce bacterial load, while lactoferrin inhibits microbial growth by binding iron, a nutrient required for microbial proliferation (Nawaz et al., 2022). Immunoglobulin A contributes to mucosal immunity by neutralizing pathogens, and antimicrobial peptides such as defensins further strengthen microbial inhibition (Zhou et al., 2025). These properties support the hypothesis that saliva can function as a local antimicrobial fluid during licking, grooming, and wound contact.

Wound-Healing Components in Saliva

In addition to antimicrobial activity, saliva contains bioactive molecules involved in wound repair. Histatin peptides, particularly histatin-1, have been shown to promote cell migration and tissue repair (Pan et al., 2021). Salivary growth-related factors may also support epithelial regeneration and angiogenesis (Yılmaz Şaştım et al., 2021). Evidence from oral and mucosal wound-healing research indicates that saliva-associated factors can influence repair dynamics, although species-specific effects in animal skin wounds require direct experimental testing (Chuhuaicura et al., 2025). Animals that lick wounds may therefore receive both antimicrobial exposure and biochemical signals associated with tissue repair. The proposed causal pathways are summarized in Table 1.

Table 1. Proposed integrated biological self-cleaning mechanisms in animals.

| Mechanism | Biological Source | Active Components | Function | Scientific Significance |
|---------------------------|--------------------------|-----------------------------------|---------------------------------|--------------------------------|
| Saliva-mediated cleaning | Salivary glands | Histatins, lysozyme, lactoferrin | Antimicrobial and wound healing | Innate immune defense |
| Self-grooming behavior | Behavioral response | Mechanical cleaning + saliva | Removal of dirt and parasites | Natural hygiene maintenance |
| Tear antimicrobial system | Lacrimal glands | Lysozyme, immunoglobulins | Ocular surface protection | Eye-cleaning mechanism |
| Skin protective barrier | Sebaceous glands | Lipids and antimicrobial peptides | Skin protection | Infection prevention |
| Overnight grooming | Resting behavior | Combined biological mechanisms | Self-cleaning during rest | Morning cleanliness phenomenon |

Grooming Behavior as a Natural Self-Cleaning Mechanism

Self-Grooming in Animals

Self-grooming is a widespread behavior with multiple biological functions. Cats spend a substantial portion of their active time grooming, and dogs, rodents, livestock, and many other animals also engage in licking or mechanical grooming. Grooming removes dirt and debris, reduces ectoparasite burden, maintains fur and skin condition, distributes protective skin oils, and can contribute to thermoregulation. Through these mechanical and biochemical effects, grooming may reduce microbial load on the skin and fur.

Grooming During Rest and Overnight Periods

Animals commonly groom when resting, undisturbed, or preparing to sleep. Overnight or other extended rest periods may provide sufficient time for repeated grooming and redistribution of saliva and skin secretions. This pattern may help explain why animals exposed to dirt or minor superficial contamination can appear cleaner after a rest interval. During grooming, saliva is spread over the fur and skin, where it may assist in contaminant removal and local microbial suppression (Webber, 2023).

Tear Secretions and Eye-Cleaning Mechanisms

Antimicrobial Components in Tears

Tear fluid plays an essential role in ocular surface defense. Tears contain antimicrobial proteins, including lysozyme, lactoferrin, and immunoglobulins, that protect the eye from pathogens and help remove debris (Chang & Purt, 2021; Eshac et al., 2021). Continuous tear production maintains ocular moisture and flushes foreign particles from the

eye surface. This system contributes to natural ocular hygiene in animals (Soimala, 2020).

Morning Ocular Deposits and Natural Cleaning

Animals may show small periocular deposits after sleep. These deposits are formed from dried tear secretions mixed with debris, dust, and microbial particles. Rather than indicating an active disease process in every case, mild deposits may reflect normal tear-mediated clearance of ocular contaminants. Persistent, excessive, colored, or painful discharge, however, should be interpreted clinically rather than as routine self-cleaning. The natural debris-clearance role of tears is consistent with broader evidence linking dust exposure and ocular irritation (Hashmi et al., 2023).

Skin Secretions and Protective Barrier

Sebaceous glands in animal skin produce lipid-rich secretions that help maintain fur condition, support barrier function, and modulate the skin surface environment. These secretions contribute to microbial balance and prevent excessive dryness. When combined with grooming, sebaceous lipids and associated antimicrobial peptides may enhance natural hygiene by distributing protective material across the skin and coat (Igielska-Kalwat et al., 2025; Mosca et al., 2025).

Integrated Natural Hygiene Mechanism in Animals

The proposed model integrates three mutually reinforcing pathways: saliva-mediated grooming, tear antimicrobial activity, and skin protective secretions. Together, these pathways may support self-cleaning, wound repair, reduced infection risk, and maintenance

of skin and fur health. Figure 1 maps the proposed sequence from saliva and grooming through tear antimicrobial action and skin secretions to three outcomes: reduced microbial load, wound healing, and a cleaner external appearance. This framework is

consistent with broader interest in natural antimicrobial compounds and salivary biomarkers in veterinary medicine (Jadav et al., 2025; Vercelli et al., 2025).




Figure 1: Proposed Natural and Self-Cleaning Mechanism in Animals

Hypothesis

It is hypothesized that animals maintain hygiene and promote superficial wound repair through an integrated biological system involving saliva-mediated grooming, tear antimicrobial secretions, and skin protective mechanisms. These processes may collectively reduce microbial load, remove debris, support barrier maintenance, and facilitate tissue repair, particularly during rest periods. The model does not imply that licking or grooming is always clinically beneficial; excessive licking can delay healing or worsen lesions in some conditions. Rather, the hypothesis proposes that normal grooming, saliva chemistry, tear defense, and sebaceous barrier functions together represent an evolutionarily conserved innate strategy for maintaining health and reducing infection risk.

Future Research Directions

Future research should compare saliva composition across species, quantify antimicrobial activity in animal saliva, evaluate changes in microbial load before and after grooming, and document grooming

patterns during rest and overnight periods. Experimental wound models could test whether saliva-derived peptides, enzymes, or growth-related molecules improve healing under controlled conditions. Studies should also distinguish adaptive grooming from excessive licking or pathological self-trauma. Such work may clarify the biological boundaries of the proposed model and support the development of bio-inspired wound-healing and antimicrobial strategies.

Conclusion

Animals possess natural behavioral, biochemical, and physiological mechanisms that may support hygiene maintenance and superficial wound repair. Saliva-mediated grooming, tear antimicrobial activity, and skin protective secretions together form a plausible integrated innate defense system. Although the hypothesis requires direct experimental validation, understanding these natural biological systems may inform veterinary hygiene, wound-care research, and the design of bio-inspired therapeutic strategies for infection control and tissue repair.

Declarations

Ethics Approval and Consent to Participate

Not applicable. This hypothesis manuscript did not involve live animals, human participants, or primary data requiring ethical approval.

Consent for Publication

Not applicable. The manuscript does not contain identifiable individual data.

Availability of Data and Materials

No datasets were generated or analyzed for this manuscript. All sources used to develop the hypothesis are cited in the reference list.

Competing Interests

The authors declare that there are no competing interests.

Funding

The authors received no specific funding for the preparation of this manuscript.

Acknowledgements

The authors thank the Nuclear Institute for Food and Agriculture (NIFA), Peshawar, Pakistan Atomic Energy Commission (PAEC), for logistical support.

Author Contributions

TA conceptualized the hypothesis, reviewed the literature, collected relevant material, prepared the visualization, and wrote and revised the manuscript. MA reviewed and edited the manuscript and contributed to conceptual refinement.

References

- Chang, A. Y., & Purt, B. (2021). Biochemistry, tear film. In StatPearls [Internet]. Treasure Island, FL: StatPearls Publishing.
- Chuhuaicura, P., Rodríguez-Niklitschek, C., Oporto, G. H., & Salazar, L. A. (2025). Distinct molecular mechanisms in oral mucosal wound healing: translational insights and future directions. *International Journal of Molecular Sciences*, 26(21), 10660.
- Eshac, Y., Redfern, R. L., & Aakalu, V. K. (2021). The role of endogenous antimicrobial peptides in modulating innate immunity of the ocular surface in dry eye diseases. *International Journal of Molecular Sciences*, 22(2), 721.
- Haick, H., & Zheng, Y. (2025). Self-cleaning function. In *Nature-Inspired Sensors* (pp. 171-179). Elsevier.
- Hart, B. L., & Hart, L. A. (2018). How mammals stay healthy in nature: the evolution of behaviours to avoid parasites and pathogens. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1751), 20170205.
- Hashmi, M. Z., Qadeer, Q., Farooq, U., & Pongpiachan, S. (2023). Dust and eye afflictions. In *Dust and Health: Challenges and Solutions* (pp. 79-92). Cham: Springer International Publishing.
- Igielska-Kalwat, J., Kilian-Pięta, E., & Siedlecki, M. (2025). The effect of a preparation derived from fish skin on the change in the fatty acid composition of sebum in individuals with demodicosis. *Lipids*, 60(6), 423-433.
- Jaday, M. M., Karmbir, & Joshi, V. (2025). Saliva as alternative diagnostic biofluid for non-invasive health and disease appraisal in domestic and wild animals: A review of salivary biomarkers and sialochemistry. *Acta Veterinaria Hungarica*, 73(4), 264-276.
- Maddu, N. (2019). Functions of saliva. In *Saliva and Salivary Diagnostics*. IntechOpen.
- Mosca, S., Ottaviani, M., Briganti, S., Di Nardo, A., & Flori, E. (2025). The sebaceous gland: a key player in the balance between homeostasis and inflammatory skin diseases. *Cells*, 14(10), 747.
- Mudgil, P. (2022). Antimicrobial tear lipids in the ocular surface defense. *Frontiers in Cellular and Infection Microbiology*, 12, 866900.
- Nawaz, N., Wen, S., Wang, F., Nawaz, S., Raza, J., Iftikhar, M., & Usman, M. (2022). Lysozyme and its application as antibacterial agent in food industry. *Molecules*, 27(19), 6305.
- Pan, L., Zhang, X., & Gao, Q. (2021). Effects and mechanisms of histatins as novel skin wound-healing agents. *Journal of Tissue Viability*, 30(2), 190-195.
- Scavizzi, F., Galligioni, V., Vasina, V., & Raspa, M. (2022). Animal health management and hygiene. In *Practical Handbook on the 3Rs in the Context of the Directive 2010/63/EU* (pp. 151-179). Academic Press.
- Soimala, T. (2020). Hygiene in Ophthalmic Surgery and Bacterial Resistance on the Ocular Surface of Animals (Doctoral dissertation).
- Tamai, R., & Kiyoura, Y. (2025). Candida infections: The role of saliva in oral health: A narrative review. *Microorganisms*, 13(4), 717.
- Tynes, V. V. (2025). Grooming behavior. In *Feline Behavior and Welfare* (pp. 83-89). Academic Press.

18. Vercelli, C., Amadori, M., Gambino, G., Danieli, D., Crimi, S., & Re, G. (2025). Natural antimicrobial compounds in veterinary medicine: Focus on companion animals. *Applied Sciences*, 15(23), 12388.
19. Vila, T., Rizk, A. M., Sultan, A. S., & Jabra-Rizk, M. A. (2019). The power of saliva: Antimicrobial and beyond. *PLoS Pathogens*, 15(11), e1008058.
20. Webber, R. (2023). Comprehensive ethogram for the American lobster (*Homarus americanus*).
21. Yılmaz Şaştım, Ç., Gürsoy, M., Könönen, E., Kasurinen, A., Norvio, S., Gürsoy, U. K., & Doğan, B. (2021). Salivary and serum markers of angiogenesis in periodontitis in relation to smoking. *Clinical Oral Investigations*, 25(3), 1117-1126.
22. Zhou, X., Wu, Y., Zhu, Z., Lu, C., Zhang, C., Zeng, L., & Zhou, F. (2025). Mucosal immune response in biology, disease prevention and treatment. *Signal Transduction and Targeted Therapy*, 10(1), 7.